

New records of mites (Arachnida: Acari) from Sable Island, Nova Scotia, Canada

Christopher G. Majka¹

Nova Scotia Museum, 1747 Summer Street, Halifax, Nova Scotia, Canada B3H 3A6

Valerie Behan-Pelletier

Agriculture and Agri-Food Canada, 960 Carling Avenue, Ottawa, Ontario, Canada K1A 0C6

Daria Bajerlein, Jerzy Błoszyk

Adam Mickiewicz University, Department of Animal Taxonomy and Ecology, Umultowska 89, Poznan 61-614, Poland

Gerald W. Krantz

Oregon State University, Department of Zoology, Cordley 3029, Corvallis, Oregon 97331-2914, United States of America

Zoe Lucas

Nova Scotia Museum, 1747 Summer Street, Halifax, Nova Scotia, Canada B3H 2G5

Barry OConnor

University of Michigan, Museum of Zoology, 1109 Geddes Avenue, Ann Arbor, Michigan 48109-1079, United States of America

Ian M. Smith

Agriculture and Agri-Food Canada, 960 Carling Avenue, Ottawa, Ontario, Canada K1A 0C6

Abstract—The first investigations of the mite fauna of Sable Island, Nova Scotia, Canada, are reported. Fourteen species have been found. *Uropoda orbicularis* (Müller) (Uropodidae) and *Scarabaspis inexpectatus* (Oudemans) (Eviphididae) are newly recorded for North America, *Macrocheles nemerdarius* Krantz and Whitaker (Macrochelidae) is newly recorded for Canada, and *Trichoribates striatus* Hammer (Ceratozetidae) is recorded for the first time south of the subarctic zone. Colonization, dispersal, and the zoogeographic origins of the fauna are discussed in the context of the biological, geological, and human history of the island.

Résumé—Cet article présente les premières investigations de la faune acarienne de l'île de Sable en Nouvelle-Écosse au Canada. On a trouvé 14 espèces. On signale le *Uropoda orbicularis* (Müller) (Uropodidae) et le *Scarabaspis inexpectatus* (Oudemans) (Eviphididae) pour la première fois en Amérique du Nord, le *Macrocheles nemerdarius* Krantz et Whitaker (Macrochelidae) pour la première fois au Canada et le *Trichoribates striatus* Hammer (Ceratozetidae) pour la première fois au sud de la zone subarctique. La colonisation, la dispersion et les origines zoogéographiques de cette faune sont abordées dans le contexte de l'histoire biologique, géologique et humaine de l'île.


Introduction

Sable Island (44°N, 60°W) is a 45 km long sand bar located near the edge of the continental shelf, 160 km from the nearest point of land

(Fig. 1). Rising no more than 25 m in height, it is covered by heathland and a variety of grasses. In the widest areas (approximately 1.5 km) there are shallow brackish and freshwater ponds. The island was ice-covered during the last phase of

Received 13 December 2006. Accepted 6 April 2007.

¹Corresponding author (e-mail: c.majka@ns.sympatico.ca).

Fig. 1. Map of Atlantic Canada and the continental shelf, indicating the position of Sable Island.

the Wisconsinian glaciation and became ice-free approximately 14 500 years ago (King 2001). Humans visited and lived on the island sporadically during the 1600s and 1700s, and it has been continuously inhabited since 1801 (Campbell 1974).

Because of its geographical situation, environmental vulnerability, long history of human activities, and unique fauna, Sable Island has been the focus of considerable scientific interest. Several endemic species have been described from the island, including *Koerneria mulveyi* Ebsary (Nematoda), *Aequipecten irradians sablensis* Clarke (Mollusca: Bivalvia), *Tricholochmaea sablensis* (Brown) (Coleoptera: Chrysomelidae), *Orygia leucostigma sablensis* Neil (Lepidoptera: Lymantriidae), *Agrotis arenarius* Neil (Lepidoptera: Noctuidae), a still undescribed species of *Papaipema* Smith (Lepidoptera: Noctuidae), as well as the well-known Ipswich sparrow (*Passerculus sandwichensis princeps* Maynard) (Aves) (Wright 1989).

Biological investigations of the island began in the middle of the nineteenth century (Gilpin 1858). Contemporary studies of its arthropod fauna began with Howden *et al.* (1970). Wright (1989) published a comprehensive treatment of the island's fauna (reporting 561

arthropod species of which 486 are insects). He recorded 28 species of spiders (Arachnida: Araneae) and 2 species of opilionids (Arachnida: Opiliones), but no Acari have been reported. This paper reports the results of preliminary investigations of the island's acarine fauna.

Materials and methods

Since 2004, Z. Lucas has been collecting insects on Sable Island. Methods have included sweep-netting, dip-netting, pitfall-trapping, and Berlese-funnel extractions, but collection has relied primarily on manual searching. Specimens of Acari have been collected incidentally by means of all these techniques and voucher specimens deposited with the following institutions:

- NSMC Nova Scotia Museum, Halifax, Nova Scotia, Canada
- OSU Oregon State University, Corvallis, Oregon, United States of America
- UMMZ University of Michigan Museum of Zoology, Ann Arbor, Michigan, United States of America

Examination of specimens at the following institutions has yielded additional records:

- CNC Canadian National Collection of Insects, Arachnids, and Nematodes, Ottawa, Ontario, Canada
 NSMC Nova Scotia Museum, Halifax, Nova Scotia, Canada
 SUNY State University of New York, Albany, New York, United States of America

Classification follows Krantz and Walter (2008).

Results

Fourteen species of mites have been found on Sable Island. Table 1 lists them and summarizes information on their bionomics and distribution. Brief notes on the Sable Island fauna follow.

Acari: Parasitiformes: Mesostigmata: Monogynaspida

Uropodina: Uropodidae

Uropoda orbicularis (Müller, 1776)

Deutonymphs were found on the scarab beetles *Aphodius fimetarius* (L.), *Aphodius subterraneus* (L.), and *Onthophagus nuchicornis* (L.), which are associated with horse dung (NSMC). *Aegialia arenaria* (F.) (Scarabaeidae), *Cercyon haemorrhoidalis* (F.), *Cercyon pygmaeus* (Illiger), and *Cryptopleurum minutum* (F.) (Hydrophilidae), all Palearctic beetles also associated with horse dung on Sable Island, were not found to host *U. orbicularis*, although *C. haemorrhoidalis* has been recorded as a host in Europe (Bajerlein and Przewoźny 2005). *Aphodius* spp. and *O. nuchicornis* are known hosts of *U. orbicularis* in Europe (Bajerlein and Błoszyk 2004). Although specimens of this species from Alberta and Newfoundland exist (CNC; E.E. Lindquist, personal communication), this is the first published report of its occurrence in North America.

Gamasina: Parasitidae

Parasitus fimetorum (Berlese, 1903)

A single deutonymph was recovered from horse manure (OSU). The species is common in small-mammal nests, rotting vegetation, dung, and compost. Specimens have also been taken from the nests of birds, and occasionally from leaf litter or open grasslands (Hyatt 1980;

Hennessey and Farrier 1988). Deutonymphs of *P. fimetorum* may be phoretic on a wide range of arthropods, including dung and carrion beetles (*Aphodius*, *Geotrupes*, *Hister*, *Nicrophorus* spp.) and bumblebees (Holzmann 1969; Richards and Richards 1976).

Gamasina: Eviphididae

Scarabaspis inexpectatus (Oudemans, 1903)

Deutonymphs were taken from the gular region of the dung beetle *O. nuchicornis* (OSU). The species has been collected from cattle dung in Europe (Karg 1993). These collections represent the first records of *S. inexpectatus* in North America. Its occurrence on Sable Island is noteworthy also in light of records from other insular sites in Europe (Borkum Island in the East Frisian Islands chain, Isles of Scilly) (Oudemans 1903; Evans 1957).

Gamasina: Macrochelidae

Macrocheles perglaber Filipponi and Peggazano, 1962

Found associated with horse dung and with dung beetles (*Aphodius* spp.) (OSU). A common Holarctic phoretic species found in domestic animal dung, where it preys on fly eggs and larvae (Filipponi 1959; Krauss 1970; Glida *et al.* 2003). It is considered a potentially useful biological agent for controlling muscine pests (Krantz 1983).

Macrocheles nemerdarius Krantz and Whitaker, 1988

Collected from beneath dead seabirds (OSU). The species was described from rodent nests in Maryland and Florida (Krantz and Whitaker 1988), where it was probably feeding on nematodes and other nidicolous microfaunal species. The female specimens from Sable Island deviate from the original description in having somewhat more robust idiosomatic setae than those found in type specimens. Prolonged isolation of the Sable Island population may have contributed to the expression of this anomaly. However, the widely disparate distributions and substrates of the Sable Island collections compared with those of the type collections suggest that males of this insular population should be sought to confirm the species' identity.

Table 1. Acari found on Sable Island, Nova Scotia, Canada.

Species	Sable Island bionomics	Preferred habitats	Biology and ecology	Known distribution
Superorder: Parasitiformes				
Order: Mesostigmata				
Suborder Monogynaspida				
Cohort Uropodina				
Family Uropodidae				
<i>Uropoda orbicularis</i> *	Phoretic on <i>Aphodius</i> spp. and <i>Onthophagus nuchicornis</i> on horse dung	Dung environments	Feeds on fungal hyphae; phoretic on dung beetles	Widely distributed in Europe; adventive in North America
Cohort Gamasina				
Family Parasitidae				
<i>Parasitus fimetorum</i> *	Horse manure	Decomposing environments	Predaceous; phoretic on many arthropods, including dung and carrion insects	Widely distributed in Europe; adventive in North America
Family Eviphididae				
<i>Scarabaspis inexpectatus</i> *	Phoretic on <i>O. nuchicornis</i> on horse manure	Dung environments	Predaceous on eggs and larvae of Diptera associated with dung and dung beetles	Insular/coastal in Europe; newly recorded in North America
Family Macrochelidae				
<i>Macrocheles perglaber</i> †	Phoretic on <i>Aphodius</i> spp. on horse manure	Dung environments	Predaceous on eggs and larvae of Diptera on dung and carrion; phoretic on dung beetles	Widely distributed Holarctic species
<i>Macrocheles nemerdarius</i>	Dead seabirds	Rodent nests	Predaceous on nematodes	Southeastern USA
Family Ascidae				
<i>Chetrosetus curtipes</i> *	Pond margins	Swamps, marshes, ponds	Predaceous	Widely distributed in Europe; adventive in North America
Order: Trombidiformes				
Suborder Prostigmata				
Cohort Parasitengonina				
Family Erythraeidae				

Table 1 (concluded).

Species	Sable Island bionomics	Preferred habitats	Biology and ecology	Known distribution
<i>Leptus</i> species A	Small brackish ponds	Aquatic and semi-aquatic	Parasitic on insects and arachnids	Unknown
Family Pionidae				
<i>Piona</i> sp. near <i>rotunda</i>	Small brackish ponds	Aquatic and in wetlands	Parasitic on chironomids	Widely distributed in North and Central America
Superorder: Acariformes				
Order: Sarcotiformes				
Suborder Oribatida				
Cohort Nothrina				
Family Nothridae				
<i>Trhypochthoniellus setosus canadensis</i>	Beach drift	Aquatic and semi-aquatic	Particle-feeding mycophage	Alaska, northern Canada, and the Maritime Provinces
Family Camisiidae				
<i>Platynothrus punctatus</i> [†]	Small brackish ponds	Aquatic and semi-aquatic	Particle-feeding saprophage and mycophage	Widely distributed Holarctic species
Cohort Brachypylina				
Family Ceratozetidae				
<i>Trichoribates striatus</i>	Found on oily porpoise skull	Decomposing environments	Particle-feeding saprophage and mycophage; possibly an opportunistic predator	Subarctic (Alaska, Yukon, and northern Manitoba)
Family Hydrozetidae				
<i>Hydrozetes</i> species A	Small brackish ponds	Aquatic	Surface-grazing microphage	Unknown
Cohort Astigmatina				
Family Histio stomatidae				
<i>Pelzneria</i> species A	On <i>Creophilus maxillosus</i> and <i>Thanatophilus lapponicus</i> in decomposing environments	Decomposing and dung environments	Filter-feeding microphage phoretic on Silphidae and Staphylinidae	Widely distributed?
<i>Spinanoetus</i> species A	On <i>C. maxillosus</i> and <i>T. lapponicus</i> in decomposing environments	Decomposing and dung environments	Filter-feeding microphage phoretic on Silphidae and Staphylinidae	Widely distributed?

*Palearctic species.

†Holarctic species.

Gamasina: Ascidae***Cheiroseius (Posttrematus) curtipes* (Halbert, 1923)**

A single female was collected from the muddy edge of a coastal pond on 26–28 August 1979 by Eric Mills (OSU). Like most other platyseine ascids, *C. (P.) curtipes* is commonly encountered in humus and moss from damp habitats such as swamps, marshes, wetlands, and watercourse banks in Europe and North and Central America (Schweizer 1949; Karg 1993). Its feeding habits are unknown, although the cheliceral morphology suggests that it is predatory (Lindquist and Evans 1965).

**Trombidiformes: Prostigmata:
Parasitengonina****Parasitengonina: Erythraeidae*****Leptus* species A**

Specimens of an undescribed species in the genus *Leptus* were collected in association with small brackish ponds (NSMC and CNC). Post-larval instars of the worldwide genus *Leptus* are predators in litter and herbaceous plant habitats (Walter *et al.* 2008). Larvae parasitize a wide range of arthropods including insects and arachnids, providing regular opportunities for the mites to disperse on their hosts. There appear to be numerous species of *Leptus* in North America, including some that are morphologically similar to species living in Eurasia. Species identities and nomenclature remain uncertain pending a revision of the genus for the Holarctic region.

Parasitengonina: Pionidae***Piona* sp. near *rotunda* (Kramer, 1979)**

A number of specimens of this water mite were collected by Paul Catling in 1981 in a brackish pond (CNC). Adults and deutonymphs of the worldwide genus *Piona* are predators of cladocerans (Branchiopoda: Cladocera) and other small arthropods in standing-water habitats (Cook 1974; Smith *et al.* 2001). Larval *Piona* spp. are typically ectoparasites of adult chironomids (Diptera: Chironomidae), which provide the primary dispersal mechanism for the mites as well as the nourishment for their developing larvae (Smith 1976; Smith and Oliver 1986). Populations of the *Piona rotunda* species complex are widely distributed in ponds

and lakes throughout the Holarctic region (Cook 1960; Viets 1987), and although several species names are available, none can be used with certainty until the group is revised.

**Acariformes: Sarcoptiformes:
Oribatida****Nothrina: Nothridae*****Trhypochthoniellus setosus canadensis*****Hammer, 1952**

J.E.H. Martin collected large numbers on 8 July 1965 in beach drift (CNC and SUNY). It has a subarctic and boreal distribution throughout the Nearctic region and has been recorded from Alaska, Yukon, Northwest Territories, Nunavut, Alberta, New Brunswick, and Nova Scotia (Marshall *et al.* 1987; Behan-Pelletier and Eamer 2004). It is among the few oribatid mites found in aquatic and semi-aquatic habitats in Canada. *Trhypochthoniellus* spp. can complete their life cycle while submerged and are truly aquatic (Seniczak *et al.* 1998). *Trhypochthoniellus s. canadensis* is a particle-feeding mycophage involved in decomposition processes.

Nothrina: Camisiidae***Platynothrus punctatus* (L. Koch, 1879)**

Collected by Eric Mills from the margins of ponds on 26–28 August 1979 (CNC). Widely distributed throughout the Holarctic region and in Canada known from the Yukon, Northwest Territories, Nunavut, Manitoba, and Newfoundland (Marshall *et al.* 1987; Behan-Pelletier and Eamer 2004). It is also among the few oribatid mites found in aquatic and semi-aquatic habitats.

Brachypylina: Ceratozetidae***Trichoribates striatus* Hammer, 1952**

Collected by Z. Lucas in large numbers on 23 July 2004, 30–31 August 2004, and 1–5 September 2004 on an oily porpoise skull (NSMC and CNC). This species has a subarctic distribution and is known from the Kenai Peninsula, Alaska, the Yukon, and Churchill, Manitoba. Most species of *Trichoribates* are particle-feeding saprophages and mycophages, possibly preying opportunistically on nematodes and other microfaunal species. They are also scavengers on small dead arthropods (necrophages).

These specimens represent the first records of this species south of the subarctic zone.

Brachypylina: Hydrozetidae

***Hydrozetes* species A**

Specimens of an undescribed species of *Hydrozetes* were collected by Eric Mills on 26–28 August 1979 from pond margins (CNC). The genus *Hydrozetes* is in need of revision. Species are found throughout the world in aquatic situations, generally in association with plants (Krantz and Baker 1982). Among Oribatida only adult *Hydrozetes* spp. exhibit levitation, *i.e.*, move upwards in the water column because of an air bubble that forms in the mid-gut when they are agitated, when the light changes from high to low intensity (Newell 1945), or when adults are dislodged (Fernandez and Athias-Binche 1986).

Astigmatina: Histiosomatidae

***Pelzneria* species A**

Several specimens of this undescribed species were collected as deutonymphs phoretic on the beetles *Creophilus maxillosus* (L.) (Coleoptera: Staphylinidae) and *Thanatophilus lapponicus* (Herbst) (Coleoptera: Silphidae) associated with dead seabirds (UMMZ). Species of *Pelzneria* are restricted to carrion habitats, where they can be found inhabiting carcasses in association with beetles of the genus *Nicrophorus* (Coleoptera: Silphidae). They are filter-feeders on microorganisms. Though both *Pelzneria* and *Spinanoetus* (see below) have been studied in Europe (Scheucher 1957), they have not been formally examined in North America. B. OConnor has collected *Pelzneria* species A from *Nicrophorus* sp. in northern lower Michigan.

***Spinanoetus* species A**

Specimens of this undescribed species were collected as deutonymphs phoretic on *C. maxillosus* in horse manure and on *T. lapponicus* on dead seabirds (UMMZ). Species of *Spinanoetus* typically inhabit vertebrate carrion but have not been studied outside Europe. Feeding stages filter microorganisms from fluid with their highly modified chelicerae. B. OConnor has collected all stages of this species from deer carcasses in Michigan and found the deutonymphs phoretic on *C. maxillosus* and on species of *Oiceoptoma* and *Necrodes* (Coleoptera: Silphidae) in Michigan, New York, and Pennsylvania.

Discussion

Given the remote location of Sable Island, the origins of its fauna are of interest. Different components of the mite fauna may have employed several modes of dispersal and colonization.

Dispersal via postglacial continental-shelf bridges

Howden *et al.* (1970), Wright (1989), and Klimaszewski *et al.* (2006) have all discussed the possibility that postglacial emergent island chains served as conduits for species from refugia on George's Bank to Sable Island. Such pathways of colonization are well established in the case of the coastal-plain flora of southwestern Nova Scotia (Keddy and Wisheu 1989). Mites such as *M. nemerdarius*, *T. striatus*, *Pelzneria* species A, and *Spinanoetus* species A could have been transported to Sable Island via beach-drift materials, wind, water, or sea ice, or over existing land bridges. *Camisia anomia* Colloff, an oribatid in the cohort Nothrina, tolerated experimental submersion in seawater for 14 days, with a survival rate of 75% (Coulson *et al.* 2002). The authors noted that this length of time is sufficient to permit ocean-current transport for approximately 700 km.

Parasitic mites such as *Piona* sp. near *rotunda* and *Leptus* species A could have been transported in association with their hosts.

It is noteworthy that subarctic mites such as *T. striatus* and *T. s. canadensis* are present on Sable Island. Miller and Elias (2000) have established from postglacial paleontological evidence that between 13 760 and 12 730 years BP, beetle assemblages in Nova Scotia consisted of tundra and tree-line species. These disappear in younger sediments and are replaced by typical boreal assemblages that flourished as the climate warmed and trees began to appear. Subarctic mites could have arrived on Sable Island via continental-shelf land bridges behind the retreating glaciers and might represent a surviving relict of the former subarctic fauna of the Maritime Provinces.

Bird-mediated colonization

Alternatively, oribatids may have been transported to Sable Island from the mainland in association with birds. Krivolutsky and Lebedeva (2004a, 2004b) have collected species of *Trichoribates* from the feathers of both passeriform

and non-passeriform birds in Russia, and *Trhypochthoniellus setosus* (Willmann) was taken from the feathers of a mallard. Those authors suggested that the mites feed on fungi associated with the feathers and that transport on birds may be an important means of oribatid dispersal to islands. Avian transport may be particularly relevant in the case of aquatic and semi-aquatic Oribatida (*T. s. canadensis*, *P. punctatus*, and *Hydrozetes* species A) found on Sable Island. Fifteen species of birds regularly breed on Sable Island, including four ducks, three shorebirds, two gulls, and three terns (Lucas 2002). These species frequent aquatic environments and could serve as a conduit for mite dispersal.

Introduced taxa and human-mediated dispersal

Four introduced Palearctic species of mites are found on Sable Island: *U. orbicularis*, *P. fimetorum*, *S. inexpectatus*, and *C. (P.) curtipes*. The Holarctic *M. perglaber* may also be an introduced species. All are associated with dung environments and are phoretic on dung beetles. All dung-associated beetles found on Sable Island are introduced Palearctic species. There are no native terrestrial mammals on Sable Island and the horse (*Equus caballus* L.) is the only introduced mammal that persists. It therefore seems probable that dung beetles were inadvertently introduced in association with the importation of livestock, which commenced in 1553 when Portuguese fisherman landed cattle and pigs to serve as a source of fresh meat (Elstracke 1625). Forty-four species of Coleoptera (Howden *et al.* 1970; Wright 1989) and 69 species of vascular plants on Sable Island are introduced (Catling *et al.* 1984).

Insular evolution

Also of interest are the collections of *M. nemerdarius*, whose modification of the substrate and idiosomatic setal anomaly (see above) suggest that the Sable Island population has diverged from continental mainland populations in response to the special environmental circumstances of this isolated offshore location.

Conclusion

This account clearly represents but a first step towards identifying and cataloguing the mite fauna of Sable Island. There is every indication that more comprehensive collecting

efforts will yield additional species, as well as further information regarding the origins and bionomics of the Sable Island acarine fauna.

What is the size of this fauna? Exact parallels are difficult to find; however, on the sub-Antarctic Prince Edward Islands (46.5°S, 37.5°E), Marshall *et al.* (1999) reported 60 species of mites, while on Heard Island (53°S, 73.5°E) Marshall and Chown (2002) reported 45 species. Although there are differences between these volcanic islands and Sable Island (44°N, 60°W) in terms of size, geology, geological history, distance from nearest land, *etc.*, they do lie at similar latitudes, have similar climates, are relatively remote, provide a very limited range of habitats, and have a combination of native and introduced faunas. Therefore, those numbers may serve as an indicator of the scale of the Sable Island acarine fauna.

Acknowledgements

We are grateful to Evert E. Lindquist (Agriculture and Agri-food Canada), Eric Mills (Dalhousie University), Roy Norton (State University of New York), Heather Proctor and David Walter (University of Alberta), and Gerry Forbes and the staff of the Sable Island Station (Meteorological Service of Canada) for their assistance. Two anonymous reviewers read a previous version of the manuscript and made many constructive suggestions. Many thanks are extended to Meghan Cameron for her translation assistance. Z. Lucas and C.G. Majka thank their colleagues at the Nova Scotia Museum, David Christianson, Calum Ewing, and Andrew Hebda, for continuing support and encouragement. This work has been assisted by the Board of Governors of the Nova Scotia Museum.

References

- Bajerlein, D., and Błoszyk, J. 2004. Phoresy of *Uropoda orbicularis* by beetles (Coleoptera) associated with cattle dung in Poland. *European Journal of Entomology*, **101**: 185–188.
- Bajerlein, D., and Przewoźny, M. 2005. Coprophagous hydrophilid beetles (Coleoptera: Hydrophilidae) as carriers of phoretic deutonymphs of *Uropoda orbicularis* (Acari: Mesostigmata) in Poland. *European Journal of Entomology*, **102**: 119–122.
- Behan-Pelletier, V.M., and Eamer, B. 2004. Diversity of Oribatida in Canada [online]. Available from http://www.cbif.gc.ca/spp_pages/mites/phps/index_e.php [accessed 30 March 2007].

- Campbell, L. 1974. Sable Island, fatal and fertile crescent. Lancelot Press, Halifax, Nova Scotia.
- Catling, P.M., Freedman, B., and Lucas, Z. 1984. The vegetation and phytogeography of Sable Island, Nova Scotia. *Proceedings of the Nova Scotia Institute of Science*, **34**: 181–247.
- Cook, D.R. 1960. Water mites of the genus *Piona* in the United States (Acarina: Pionidae). *Annals of the Entomological Society of America*, **53**: 35–60.
- Cook, D.R. 1974. Water mite genera and subgenera. *Memoirs of the American Entomological Institute* No. 21.
- Coulson, S.J., Hodkinson, I.D., Webb, N.R., and Harrison, J.A. 2002. Survival of terrestrial soil-dwelling arthropods on and in seawater: implications for trans-oceanic dispersal. *Functional Ecology*, **16**: 353–356.
- Elstracke, R. 1625. *Hakjulutus posthumus* or Purchas his pilgrimes: contayning a history of the world in sea voyages and lande travells by Englishmen and others. *Edited by S. Purchas*. James MacLehose and Sons, Glasgow, UK.
- Evans, G.O. 1957. On the genus *Scarabaspis* Womersley (Acarina-Mesostigmata). *Annals and Magazine of Natural History, Series 12*, **10**: 409–416.
- Fernandez, N.A., and Athias-Binche, F. 1986. Analyse démographique d'une population d'*Hydrozetes lemnae* Coggi, Acarien Oribate inféodé à la lentille d'eau *Lemna gibba* L. en Argentine. I. Méthodes et techniques, démographie d'*H. lemnae* et comparaisons avec d'autre Oribates. *Zoologische Jahrbücher. Abteilung für Systematik*, **113**: 213–228.
- Filipponi, A. 1959. Due specie riproduttivamente isolate sotto il nome di *Macrocheles glaber* (Mueller) (Acarina, Mesostigmata). *Rivista Parassitologia*, **20**: 199–212.
- Gilpin, J.B. 1858. Sable Island: its past history, present appearance, natural history, etc., etc. Wesleyan Conference Steam Press, Halifax, Nova Scotia.
- Glida, H., Bertrand, M., and Peyrusse, V. 2003. A limiting factor in the abundance of predatory phoretic mites (Acari: Macrochelidae): the seasonal abundance of their phorionts (dung beetles) in southern France. *Canadian Journal of Zoology*, **81**: 2066–2072.
- Hennessey, M.K., and Farrier, M.H. 1988. Systematic revision of thirty species of free-living, soil-inhabiting gamasine mites (Acari: Mesostigmata) of North America. *North Carolina Agricultural Research Service Technical Bulletin* No. 285.
- Holzmann, C. 1969. Die Familie Parasitidae Oudemans 1901: eine systematische Studie aus dem Jahre 1955. *Acarologie (Schriftenreihe für Vergleichende Milbenkunde)*, **13**: 3–55.
- Howden, H.F., Martin, J.E.H., Bousfield, E.L., and McAllister, D.E. 1970. Fauna of Sable Island and its zoogeographic affinities. *National Museums of Canada Publications in Zoology*, **4**: 1–45.
- Hyatt, K.H. 1980. Mites of the subfamily Parasitinae (Mesostigmata: Parasitidae) in the British Isles. *Bulletin of the British Museum (Natural History), Zoology Series*, **38**: 237–378.
- Karg, W. 1993. Raubmilben. Acari (Acarina), Milben Parasitiformes (Anactinochaeta), Cohors Gamasina Leach. *Die Tierwelt Deutschlands, Part 59*.
- Keddy, P.A., and Wisheu, I.C. 1989. Ecology, biogeography, and conservation of coastal plain plants: some general principles from the study of Nova Scotian wetlands. *Rhodora*, **91**: 72–94.
- King, E.L. 2001. A glacial origin for Sable Island: ice and sea-level fluctuations from seismic stratigraphy on Sable Island Bank, Scotian Shelf, offshore Nova Scotia. *Geological Survey of Canada, Current Research 2001-D19*.
- Klimaszewski, J., Majka, C.G., and Langor, D. 2006. Review of the North American *Tarphiota* Casey, with a description of a new seashore-inhabiting species *Atheta* species exhibiting convergent characteristics (Coleoptera: Staphylinidae: Aleocharinae). *Entomological Science*, **9**: 67–78.
- Krantz, G.W. 1983. Mites as biological control agents of dung-breeding flies, with special reference to the Macrochelidae. *In* *Biological control of pests by mites*. *Edited by M.A. Hoy, G.L. Cunningham, and L. Knutson*. University of California Special Publications No. 3304. pp. 91–98.
- Krantz, G.W., and Baker, G.T. 1982. Observations on the plastron mechanism of *Hydrozetes* sp. (Acari: Oribatida: Hydrozetidae). *Acarologia*, **23**(3): 273–277.
- Krantz, G.W., and Walter, D.E. (Editors). 2008. *A manual of acarology*. 3rd ed. Texas Tech University Press, Lubbock, Texas. In press.
- Krantz, G.W., and Whitaker, J.O., Jr. 1988. Mites of the genus *Macrocheles* (Acari: Macrochelidae) associated with small mammals in North America. *Acarologia*, **29**: 225–259.
- Krauss, W. 1970. Die europäischen Arten der Gattungen *Macrocheles* Latreille 1829 und *Geholaspis* Berlese 1918. *Acarologie (Schriftenreihe für vergleichende Milbenkunde)*, **14**: 2–53.
- Krivolutsky, D.A., and Lebedeva, N.V. 2004a. Oribatid mites (Oribatei, Acariformes) in bird feathers: non-passerines. *Acta Zoologica Lituonica*, **14**(1): 26–47.
- Krivolutsky, D.A., and Lebedeva, N.V. 2004b. Oribatid mites (Oribatei) in bird feathers: Passeriformes. *Acta Zoologica Lituonica*, **14**(2): 19–38.
- Lindquist, E.E., and Evans, G.O. 1965. Taxonomic concepts in the Ascidae, with a modified setal nomenclature for the idiosoma of the Gamasina (Acarina: Mesostigmata). *Memoirs of the Entomological Society of Canada*, **47**: 1–64.
- Lucas, Z. 2002. Birds of Sable Island [online]. Available from <http://www.greenhorsesociety.com/Birds/birds.htm> [accessed 30 March 2007].

- Marshall, D.J., and Chown, S.L. 2002. The acarine fauna of Heard Island. *Polar Biology*, **25**(9): 688–695.
- Marshall, D.J., Gremmen, N.J.M., Coetzee, L., O'Connor, B.M., Pugh, P.J.A., Theron, P.D., and Ueckermann, E.A. 1999. New records of Acari from the sub-Antarctic Prince Edward Islands. *Polar Biology*, **21**(2): 84–89.
- Marshall, V.G., Reeves, R.M., and Norton, R.A. 1987. Catalogue of the Oribatida (Acari) of continental United States and Canada. *Memoirs of the Entomological Society of Canada* No. 139.
- Miller, R.F., and Elias, S.A. 2000. Late-glacial climate in the Maritimes region, Canada, reconstructed from mutual climatic range analysis of fossil Coleoptera. *Boreas*, **29**: 79–88.
- Newell, I.M. 1945. *Hydrozetes* Berlese (Acari, Oribatidea): the occurrence of the genus in North America and the phenomenon of levitation. *Transactions Connecticut Academy of Arts and Sciences*, **36**: 253–275.
- Oudemans, A.C. 1903. Acarologische Aanteekeningen, vii. *Entomologische Berichten*, **13**: 91–92.
- Richards, L.A., and Richards, K.W. 1976. Parasitid mites associated with bumblebees in Alberta, Canada (Acarina: Parasitidae; Hymenoptera: Apidae). I. Taxonomy. *University of Kansas Science Bulletin*, **50**: 731–773.
- Scheucher, R. 1957. Systematik und Ökologie der deutschen Anoetinen. In *Beiträge zur Systematik und Ökologie mitteleuropäischer Acarina*. Vol. 1. *Edited by H.-J. Stammer*. Akademische Verlagsgesellschaft, Leipzig, Germany. pp. 233–384.
- Schweizer, J. 1949. Die Landmilben des Schweizerischen Nationalparks. *Kommission der Schweizerischen Naturforschenden Gesellschaft, Band II (NF), Teil 1: Parasitiformes Reuter 1909*. Verlag Lüdlin A.G., Liestal, Schweiz.
- Seniczak, S., Norton, R.A., and Wang, H.-F. 1998. The morphology of juvenile stages of moss mites of the family Trhypochthoniidae (Acari: Oribatida), and the taxonomic status of some genera and species. *Zoologische Anzeiger*, **237**(1): 85–95.
- Smith, I.M. 1976. A study of the systematics of the water mite family Pionidae (Prostigmata: Parasitengona). *Memoirs of the Entomological Society of Canada* No. 98.
- Smith, I.M., and Oliver, D.R. 1986. Review of parasitic associations of larval water mites (Acari: Parasitengona: Hydrachnida) with insect hosts. *The Canadian Entomologist*, **118**: 407–472.
- Smith, I.M., Cook, D.R., and Smith, B.P. 2001. Water mites and other arachnids. In *Ecology and classification of North American freshwater invertebrates*. 2nd ed. *Edited by J. Thorpe and A. Covich*. Academic Press, San Diego, California. pp. 551–659.
- Viets, K.O. 1987. Die Milben des Süßwassers (Hydrachnellae und Halacaridae [part.], Acari). 2. Katalog. *Sonderbände des Naturwissenschaftlichen Vereins in Hamburg*, **8**.
- Walter, D.E., Lindquist, E.E., Smith, I.M., Cook, D.R., and Krantz, G.W. 2008. Prostigmata, Chap. 13. In *A manual of acarology*. 3rd ed. *Edited by G.W. Krantz and D.E. Walter*. Texas Tech University Press, Lubbock, Texas. In press.
- Wright, B. 1989. The fauna of Sable Island. *Nova Scotia Museum Curatorial Report* No. 68.